



TITLE:

Generalized Alternative Theorems based on Set-Relations (Nonlinear Analysis and Convex Analysis)

AUTHOR(S):

Ogata, Yuto; Saito, Yutaka; Tanaka, Tamaki; Lee,
Gue Myung; Lee, Jae Hyoung

CITATION:

Ogata, Yuto ...[et al]. Generalized Alternative Theorems based on Set-Relations (Nonlinear Analysis and Convex Analysis). 数理解析研究所講究録 2018, 2065: 111-119

ISSUE DATE:

2018-04

URL:

<http://hdl.handle.net/2433/241910>

RIGHT:

GENERALIZED ALTERNATIVE THEOREMS BASED ON SET-RELATIONS

YUTO OGATA, YUTAKA SAITO, TAMAKI TANAKA (NIIGATA UNIVERSITY)
GUE MYUNG LEE, JAE HYOUNG LEE (PUKYONG NATIONAL UNIVERSITY)

ABSTRACT. This paper contains generalized alternative theorems via a set scalarization method based on set-relations. We introduce semidefinite system and its dual case as a supplementary result.

1. INTRODUCTION

Gordan's theorem was appeared in 1873 [2]. This theorem focuses on the geometry of finitely many vectors and the origin. This paper is originally motivated by a generalized Gordan's theorem for vector-valued functions [4] in 1986. Vector-valued functions are able to be replaced with matrices under a convex assumption. This was interestingly extended to the case of set-valued maps [7] in 1999 and [9] in 2000. They found that set-valued maps can be used in the form of alternative theorem with a similar convexity.

Their theorems based on some separation theorems require linear scalarizations such as bilinear forms. In 2005, Nishizawa, Onodsuka, and Tanaka gave generalization forms [8] by using sublinear scalarizations for vectors inspired by ones introduced in [3]. Sublinear separation enables to eliminate assumptions of theorems related to linearity and convexity.

Kuwano, Tanaka, and Yamada established sublinear scalarizations for sets [6] in 2009 by use of set-relations [5]. These functions are composed of six infimum types and six supremum types. The number of the functions for set is much larger than that for vectors which consist of just four types. This comes from the complexity of ways where we consider a set preceding another.

In this paper, we show 12 alternative theorems by using the scalarizing functions for sets, considering the set-relations between an image and a reference set. If the reference set is the origin, our theorems will be the result of the previous research. This reference set gives flexibility and let problems free themselves from the origin.

2010 *Mathematics Subject Classification.* Primary: 90C29 ; Secondary: 90C46, 49J53, 54C60.

Key words and phrases. Alternative theorem, sublinear scalarization, set-relations, set-valued map.

This work was supported by JSPS KAKENHI Grant Number 26400194.

2. PRELIMINARIES

Unless otherwise specified, we let X be a nonempty set, Y a real ordered topological vector space with \leq_C induced by a convex solid (i.e., there exists nonempty interior) cone C as follows: $x \leq_C y$ if $y - x \in C$ for $x, y \in Y$, and F a set-valued map from X to 2^Y .

Definition 2.1 (Kuroiwa, Tanaka, Ha (1997), [5]). Let $A, B \in 2^Y \setminus \{\emptyset\}$.

- (i) $A \leq_C^{(1)} B \stackrel{\text{def}}{\iff} A \subset \bigcap_{b \in B} (b - C) \Leftrightarrow B \subset \bigcap_{a \in A} (a + C)$;
- (ii) $A \leq_C^{(2)} B \stackrel{\text{def}}{\iff} A \cap \bigcap_{b \in B} (b - C) \neq \emptyset$;
- (iii) $A \leq_C^{(3)} B \stackrel{\text{def}}{\iff} B \subset (A + C)$;
- (iv) $A \leq_C^{(4)} B \stackrel{\text{def}}{\iff} \bigcap_{a \in A} (a + C) \cap B \neq \emptyset$;
- (v) $A \leq_C^{(5)} B \stackrel{\text{def}}{\iff} A \subset (B - C)$;
- (vi) $A \leq_C^{(6)} B \stackrel{\text{def}}{\iff} A \cap (B - C) \neq \emptyset \Leftrightarrow (A + C) \cap B \neq \emptyset$.

Definition 2.2 (Kuwano, Tanaka, Yamada, (2009), [6]). Let $A, B \in 2^Y \setminus \{\emptyset\}$ and $k \in \text{int}C$. For each $j = 1, \dots, 6$, scalarizing functions $I_{k,B}^{(j)}$ and $S_{k,B}^{(j)}$ from $2^Y \setminus \{\emptyset\}$ to $\mathbb{R} \cup \{\pm\infty\}$ are defined by $I_{k,B}^{(j)}(A) := \inf\{t \in \mathbb{R} \mid A \leq_C^{(j)}(tk + B)\}$, $S_{k,B}^{(j)}(A) := \sup\{t \in \mathbb{R} \mid (tk + B) \leq_C^{(j)} A\}$.

3. GENERALIZED ALTERNATIVE THEOREMS

3.1. Infimum Types.

Theorem 3.1 (Infimum type (1)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If F is compact-valued on X and V is compact, then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $F(x) \leq_{\text{int}C}^{(1)} V$;
- (ii) there exists $k \in \text{int}C$ such that $(I_{k,V}^{(1)} \circ F)(x) \geq 0$ for all $x \in X$.

Proof. [(i) \Rightarrow not (ii)] Fix $k \in \text{int}C$. Assume that (i) is consistent. Then, there exists $x \in X$ such that $y \in \bigcap_{v \in V} (v - \text{int}C)$ for all $y \in F(x)$. Since $\bigcap_{v \in V} (v - \text{int}C)$ is open, there exists $\lambda_y > 0$ such that $y + \lambda_y k \in \bigcap_{v \in V} (v - \text{int}C)$. Therefore, $y \in S_y := \bigcap_{v \in -\lambda_y k + V} (v - \text{int}C)$. This implies $\{S_y\}_{y \in F(x)}$ is a cover of $F(x)$. Since $F(x)$ is compact, there exists $\{y_1, \dots, y_n\} \subset F(x)$ such that $F(x) \subset \bigcup_{i \in I} S_{y_i}$ where $I := \{1, \dots, n\}$.

Taking $\bar{\lambda} := \min\{\lambda_{y_1}, \dots, \lambda_{y_n}\} > 0$, then $F(x) \subset \bigcup_{i \in I} \bigcap_{v \in -\lambda_{y_i} k + V} (v - \text{int}C) = \bigcap_{v \in -\bar{\lambda} k + V} (v - \text{int}C) \subset \bigcap_{v \in -\bar{\lambda} k + V} (v - C)$. Thus, $\inf\{t \in \mathbb{R} \mid F(x) \subset \bigcap_{v \in tk + V} (v - C)\} \leq -\bar{\lambda} < 0$.

[Not (i) \Rightarrow (ii)] Assume that (i) is not consistent. Then, for all $x \in X$, there exists $\bar{y} \in F(x)$ such that $\bar{y} \notin \bigcap_{v \in V} (v - \text{int}C)$. Since $\bigcap_{v \in \lambda k + V} (v - C) \subset \bigcap_{v \in V} (v - \text{int}C)$ for all

$k \in \text{int}C$ and $\lambda < 0$, then $\bar{y} \notin \bigcap_{v \in \lambda k + V} (v - C)$. Thus, $\inf\{t \in \mathbb{R} \mid F(x) \subset \bigcap_{v \in tk + V} (v - C)\}$ is non-negative. \square

Remark 3.1. We let $S := \bigcap_{v \in V} (v - C)$. Then $S + (-C) \subset S$ so that for all $x \in X$, $y \in F(x)$, and $k \in \text{int}C$, there exists $t_y > 0$ such that $y \in \bigcap_{v \in t_y k + V} (v - C)$. Thus, there exists $\bar{t} \geq 0$ such that $(I_{k,V}^{(1)} \circ F)(x) = \inf\{t \in \mathbb{R} \mid F(x) \subset \bigcap_{v \in tk + V} (v - C)\} \leq \bar{t} < +\infty$.

Theorem 3.2 (Infimum type (2)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If V is compact, then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $F(x) \leq_{\text{int}C}^{(2)} V$;
- (ii) there exists $k \in \text{int}C$ such that $(I_{k,V}^{(2)} \circ F)(x) \geq 0$ for all $x \in X$.

Proof. [(i) \Rightarrow not (ii)] Fix $k \in \text{int}C$. Assume that (i) is consistent. Then, there exist $x \in X$ and $\bar{y} \in F(x)$ such that $\bar{y} \in \bigcap_{v \in V} (v - \text{int}C)$. Since $\bigcap_{v \in V} (v - \text{int}C)$ is open, there exists $\bar{\lambda} > 0$ such that $\bar{y} + \bar{\lambda}k \in \bigcap_{v \in V} (v - \text{int}C) \subset \bigcap_{v \in V} (v - C)$. Therefore, $\bar{y} \in \bigcap_{v \in -\bar{\lambda}k + V} (v - C)$. Then, $F(x) \cap \bigcap_{v \in -\bar{\lambda}k + V} (v - C) \neq \emptyset$. Thus, $\inf\{t \in \mathbb{R} \mid F(x) \cap \bigcap_{v \in tk + V} (v - C) \neq \emptyset\} \leq -\bar{\lambda} < 0$.

[Not (i) \Rightarrow (ii)] Assume that (i) is not consistent. Then, $y \notin \bigcap_{v \in V} (v - \text{int}C)$ for all $x \in X$ and $y \in F(x)$. That means, $y \notin \bigcap_{v \in \lambda k + V} (v - C)$ for all $k \in \text{int}C$ and $\lambda < 0$. Thus, $\inf\{t \in \mathbb{R} \mid F(x) \cap \bigcap_{v \in tk + V} (v - C) \neq \emptyset\}$ is non-negative. \square

Remark 3.2. We let $S := \bigcap_{v \in V} (v - C)$ like Remark 3.1. For all $x \in X$, $y \in F(x)$, and $k \in \text{int}C$, there exists $\bar{t} > 0$ such that $y \in \bigcap_{v \in \bar{t}k + V} (v - C)$. Thus, $(I_{k,V}^{(2)} \circ F)(x) = \inf\{t \in \mathbb{R} \mid F(x) \cap \bigcap_{v \in tk + V} (v - C) \neq \emptyset\} \leq \bar{t} < +\infty$.

Theorem 3.3 (Infimum type (3)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If V is compact, then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $F(x) \leq_{\text{int}C}^{(3)} V$;
- (ii) there exists $k \in \text{int}C$ such that $(I_{k,V}^{(3)} \circ F)(x) \geq 0$ for all $x \in X$.

Proof. [(i) \Rightarrow not (ii)] Fix $k \in \text{int}C$. Assume that (i) is consistent. Then, there exists $x \in X$ such that $v \in F(x) + \text{int}C$ for all $v \in V$. Since $F(x) + \text{int}C$ is open, there exists $\lambda_v > 0$ such that $v - \lambda_v k \in F(x) + \text{int}C$. We let $S_v := \lambda_v k + F(x) + \text{int}C$ then, $\{S_v\}_{v \in V}$ is a cover of V . Since V is compact, there exists $\{v_1, \dots, v_n\} \subset V$ such that $V \subset \bigcup_{i \in I} S_{v_i}$, where $I := \{1, \dots, n\}$.

Taking $\bar{\lambda} := \min \{\lambda_{v_1}, \dots, \lambda_{v_n}\} > 0$, then $V \subset \bigcup_{i \in I} (\lambda_{v_i} k + F(x) + \text{int}C) = \bar{\lambda}k + F(x) + \text{int}C \subset \bar{\lambda}k + F(x) + C$. Thus, $\inf\{t \in \mathbb{R} \mid (tk + V) \subset (F(x) + C)\} \leq -\bar{\lambda} < 0$.

[Not (i) \Rightarrow (ii)] Assume that (i) is not consistent. Then, for all $x \in X$, there exists $\bar{v} \in V$ such that $\bar{v} \notin F(x) + \text{int}C$. Since $-\lambda k + F(x) + C \subset F(x) + \text{int}C$ for all $k \in \text{int}C$ and $\lambda < 0$, then $\bar{v} \notin -\lambda k + F(x) + C$. Thus, $\inf\{t \in \mathbb{R} \mid (tk + V) \subset (F(x) + C)\}$ is non-negative. \square

Remark 3.3. We let $S := F(x) + C$. Then $S + C \subset S$ so that for all $v \in V$ and $k \in \text{int}C$, there exists $\bar{t} > 0$ such that $\bar{t}k + v \in F(x) + C$. Thus, $(I_{k,V}^{(3)} \circ F)(x) = \inf\{t \in \mathbb{R} \mid (tk + V) \subset (F(x) + C)\} \leq \bar{t} < +\infty$.

Theorem 3.4 (Infimum type (4)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If F is compact-valued on X , then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $F(x) \leq_{\text{int}C}^{(4)} V$;
- (ii) there exists $k \in \text{int}C$ such that $(I_{k,V}^{(4)} \circ F)(x) \geq 0$ for all $x \in X$.

Proof. [(i) \Rightarrow not (ii)] Fix $k \in \text{int}C$. Assume that (i) is consistent. Then, there exist $x \in X$ and $\bar{v} \in V$ such that $\bar{v} \in \bigcap_{y \in F(x)} (y + \text{int}C)$. Since $\bigcap_{y \in F(x)} (y + \text{int}C)$ is open, there exists $\bar{\lambda} > 0$ such that $-\bar{\lambda}k + \bar{v} \in \bigcap_{y \in F(x)} (y + \text{int}C) \subset \bigcap_{y \in F(x)} (y + C)$.

Thus, $\inf\{t \in \mathbb{R} \mid (tk + V) \cap \bigcap_{y \in F(x)} (y + C) \neq \emptyset\} \leq -\bar{\lambda} < 0$.

[Not (i) \Rightarrow (ii)] Assume that (i) is not consistent. Then, $v \notin \bigcap_{y \in F(x)} (y + \text{int}C)$ for all $x \in X$ and $v \in V$. Since $\bigcap_{y \in F(x)} (-\lambda k + y + C) \subset \bigcap_{y \in F(x)} (y + \text{int}C)$ for all $k \in \text{int}C$ and $\lambda < 0$, then $v \notin \bigcap_{y \in F(x)} (-\lambda k + y + C)$.

Thus, $\inf\{t \in \mathbb{R} \mid (tk + V) \cap \bigcap_{y \in F(x)} (y + C) \neq \emptyset\}$ is non-negative. \square

Remark 3.4. We let $S := \bigcap_{y \in F(x)} (y + C)$. Then $S + C \subset S$ so that for all $v \in V$ and $k \in \text{int}C$, there exists $t_v > 0$ such that $t_v k + v \in \bigcap_{y \in F(x)} (y + C)$. Then, there exists $\bar{t} \geq 0$ such that $(\bar{t}k + V) \cap \bigcap_{y \in F(x)} (y + C) \neq \emptyset$. Thus, $(I_{k,V}^{(4)} \circ F)(x) = \inf\{t \in \mathbb{R} \mid (tk + V) \cap \bigcap_{y \in F(x)} (y + C) \neq \emptyset\} \leq \bar{t} < +\infty$.

Theorem 3.5 (Infimum type (5)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If F is compact-valued on X , then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $F(x) \leq_{\text{int}C}^{(5)} V$;
- (ii) there exists $k \in \text{int}C$ such that $(I_{k,V}^{(5)} \circ F)(x) \geq 0$ for all $x \in X$.

Proof. [(i)⇒not (ii)] Fix $k \in \text{int}C$. Assume that (i) is consistent. Then, there exists $x \in X$ such that $y \in V - \text{int}C$ for all $y \in F(x)$. Since $V - \text{int}C$ is open, there exists $\lambda_y > 0$ such that $y + \lambda_y k \in V - \text{int}C$. We let $S_y := -\lambda_y k + V - \text{int}C$, then $\{S_y\}_{y \in F(x)}$ is a cover of $F(x)$. Since $F(x)$ is compact, there exists $\{y_1, \dots, y_n\} \subset F(x)$ such that $F(x) \subset \bigcup_{i \in I} S_{y_i}$ where $I := \{1, \dots, n\}$.

Taking $\bar{\lambda} := \min\{\lambda_{y_1}, \dots, \lambda_{y_n}\} > 0$, then $F(x) \subset \bigcup_{i \in I} (-\lambda_{y_i} k + V - \text{int}C) = -\bar{\lambda}k + V - \text{int}C \subset -\bar{\lambda}k + V - C$. Thus, $\inf\{t \in \mathbb{R} \mid F(x) \subset tk + V - C\} \leq -\bar{\lambda} < 0$.

[Not (i)⇒(ii)] Assume that (i) is not consistent. Then, for all $x \in X$, there exists $\bar{y} \in F(x)$ such that $\bar{y} \notin V - \text{int}C$. Since $\lambda k + V - C \subset V - \text{int}C$ for all $k \in \text{int}C$ and $\lambda < 0$, then $\bar{y} \notin \lambda k + V - C$. Thus, $\inf\{t \in \mathbb{R} \mid F(x) \subset tk + V - C\}$ is non-negative. \square

Remark 3.5. We let $S := V - C$. Then $S + (-C) \subset S$ so that for all $x \in X, y \in F(x)$, and $k \in \text{int}C$, there exists $t_y > 0$ such that $y \in t_y k + V - C$. Then, there exists $\bar{t} \geq 0$ such that $F(x) \subset \bar{t}k + V - C$. Thus, $(I_{k,V}^{(5)} \circ F)(x) = \inf\{t \in \mathbb{R} \mid F(x) \subset (tk + V - C)\} \leq \bar{t} < +\infty$.

Theorem 3.6 (Infimum type (6)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . Then, exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $F(x) \leq_{\text{int}C}^{(6)} V$;
- (ii) there exists $k \in \text{int}C$ such that $(I_{k,V}^{(6)} \circ F)(x) \geq 0$ for all $x \in X$.

Proof. [(i)⇒not (ii)] Fix $k \in \text{int}C$. Assume that (i) is consistent. Then, there exist $x \in X$ and $\bar{y} \in F(x)$ such that $\bar{y} \in V - \text{int}C$. Thus, there exists $\bar{\lambda} > 0$ such that $\bar{y} + \bar{\lambda}k \in V - \text{int}C \subset V - C$. Then, $F(x) \cap (-\bar{\lambda}k + V - C) \neq \emptyset$. Therefore, $\inf\{t \in \mathbb{R} \mid F(x) \cap (tk + V - C) \neq \emptyset\} \leq -\bar{\lambda} < 0$.

[Not (i)⇒(ii)] Assume that (ii) is not consistent. Then, $y \in V - \text{int}C$ for all $x \in X$ and $y \in F(x)$. Since $\lambda k + V - C \subset V - \text{int}C$ for all $k \in \text{int}C$ and $\lambda < 0$, then $y \notin \lambda k + V - C$. Thus, $\inf\{t \in \mathbb{R} \mid F(x) \cap (tk + V - C) \neq \emptyset\}$ is non-negative. \square

Remark 3.6. We let $S := V - C$ like Remark 3.5. For all $x \in X, y \in F(x)$, and $k \in \text{int}C$, there exists $t_y > 0$ such that $y \in t_y k + V - C$. Then, there exists $\bar{t} \geq 0$ such that $F(x) \cap (\bar{t}k + V - C) \neq \emptyset$. Thus, $(I_{k,V}^{(6)} \circ F)(x) = \inf\{t \in \mathbb{R} \mid F(x) \cap (tk + V - C) \neq \emptyset\} \leq \bar{t} < +\infty$.

3.2. Supremum Types. We obtain supremum types conversely by replacing an ordering cone C in Infimum types with $-C$. In addition, just like six remarks described above, $(S_{k,V}^{(j)} \circ F)(x) > -\infty$ for all $x \in X$ and $j = 1, \dots, 6$.

Theorem 3.7 (Supremum type (1)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If F is compact-valued on X and V is compact, then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $V \leq_{\text{int}C}^{(1)} F(x)$;
- (ii) there exists $k \in \text{int}C$ such that $(S_{k,V}^{(1)} \circ F)(x) \leq 0$ for all $x \in X$.

Theorem 3.8 (Supremum type (2)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If V is compact, then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $V \leq_{\text{int}C}^{(2)} F(x)$;
- (ii) there exists $k \in \text{int}C$ such that $(S_{k,V}^{(2)} \circ F)(x) \leq 0$ for all $x \in X$.

Theorem 3.9 (Supremum type (3)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If V is compact, then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $V \leq_{\text{int}C}^{(3)} F(x)$;
- (ii) there exists $k \in \text{int}C$ such that $(S_{k,V}^{(3)} \circ F)(x) \leq 0$ for all $x \in X$.

Theorem 3.10 (Supremum type (4)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If F is compact-valued on X , then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $V \leq_{\text{int}C}^{(4)} F(x)$;
- (ii) there exists $k \in \text{int}C$ such that $(S_{k,V}^{(4)} \circ F)(x) \leq 0$ for all $x \in X$.

Theorem 3.11 (Supremum type (5)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . If F is compact-valued on X , then exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $V \leq_{\text{int}C}^{(5)} F(x)$;
- (ii) there exists $k \in \text{int}C$ such that $(S_{k,V}^{(5)} \circ F)(x) \leq 0$ for all $x \in X$.

Theorem 3.12 (Supremum type (6)). Let X be a nonempty set, Y a topological vector space, C a convex solid cone in Y , $F : X \rightarrow 2^Y$ a set-valued map, and V a nonempty subset in Y . Then, exactly one of the following two systems is consistent;

- (i) there exists $x \in X$ such that $V \leq_{\text{int}C}^{(6)} F(x)$;

- (ii) there exists $k \in \text{int}C$ such that $(S_{k,V}^{(6)} \circ F)(x) \leq 0$ for all $x \in X$.

4. POSITIVE SEMI-DEFINITE SYSTEM

Let S^n be the set of $n \times n$ symmetric matrices, S_+^n the set of $n \times n$ positive semidefinite symmetric matrices, S_{++}^n the set of $n \times n$ positive definite symmetric matrices, $F : X \rightarrow 2^{S^n}$ a set-valued map, and V a nonempty subset in S^n . We denote the trace of $A \in S^n$ by $\text{tr}(A) := \sum_{i=1}^n a_{ii}$. Some notions are from pp.9–10 of [1].

Theorem 4.1. Exactly one of the following two systems is consistent;

- (i) There exists $x \in X$ such that $F(x) \cap (V - S_{++}^n) \neq \emptyset$;
- (ii) There exists $K \in S_{++}^n$ such that

$$\inf\{t \in \mathbb{R} \mid F(x) \cap (tK + V - S_+^n) \neq \emptyset\} \geq 0 \text{ for all } x \in X.$$

This theorem is simply verified by $S_{++}^n = \text{int}S_+^n$.

Corollary 4.1. Let $f : \mathbb{R}^m \rightarrow S^n$ be a function. Then, exactly one of the following two systems is consistent;

- (i) There exists $x \in \mathbb{R}^m$ such that $f(x) \in -S_{++}^n$;
- (ii) There exists $B \in S_{++}^n$ such that

$$\inf\{t \in \mathbb{R} \mid f(x) \in tB - S_+^n\} \geq 0 \text{ for all } x \in \mathbb{R}^m.$$

Let $A \in S^n$ and $B \in S_{++}^n$ be given. Consider the following semidefinite optimization problem and its dual problem:

$$\begin{aligned} (\text{SDP}) \quad & \inf\{\text{tr}(-AX) \mid \text{tr}((-B)X) = -1, X \in S_+^n\}. \\ (\text{SDD}) \quad & \sup\{-t \mid -tB + S = -A, S \in S_+^n\}. \end{aligned}$$

Notice that $\inf\{t \in \mathbb{R} \mid A \in tB - S_+^n\} = -\sup\{-t \in \mathbb{R} \mid -tB + S = -A, S \in S_+^n\}$ for all $A \in S^n$ and $B \in S_{++}^n$, and we can calculate $\inf\{t \in \mathbb{R} \mid A \in tB - S_+^n\}$ with Matlab as a semidefinite optimization problem.

Lemma 4.1. Problems (SDP) and (SDD) are strictly feasible, that is, there exists $\tilde{X} \in S_{++}^n$ such that $\text{tr}((-B)\tilde{X}) = -1$, and there exist $\tilde{t} \in \mathbb{R}$ and $\tilde{S} \in S_{++}^n$ such that $-\tilde{t}B + \tilde{S} = -A$.

Proof. Let $X := \sum_{i=1}^n u_i q_i q_i^T \in S_{++}^n$ where u_i are positive and $\{q_i\}_{i=1,\dots,n}$ is a base of \mathbb{R}^n . Then, $\text{tr}(BX) = \text{tr}(B(\sum_{i=1}^n u_i q_i q_i^T)) = \sum_{i=1}^n u_i \text{tr}(Bq_i q_i^T) = \sum_{i=1}^n u_i q_i^T B q_i > 0$. Thus, $\tilde{X} := (1/\text{tr}(BX))X \in S_{++}^n$ satisfies $\text{tr}(B\tilde{X}) = 1$.

Let $B \in S_{++}^n$. If $A = \theta_{S^n}$, we have $-tB + \tilde{S} = \theta_{S^n}$ by letting $\tilde{S} := tB$ for all $t > 0$. Now we assume that $A \neq \theta_{S^n}$. Since S_{++}^n is open, there exists an open neighborhood

$U \in N(\theta_{S^n})$ such that $B + U \subset S_{++}^n$. There exists $\tilde{t} > 0$ such that $(-1/\tilde{t})A \in U$. Therefore, we can find $\tilde{S} := \tilde{t}B - A = \tilde{t}(B - (1/\tilde{t})A) \in S_{++}^n$. \square

Lemma 4.2. The optimal values of (SDP) and (SDD) coincide and their optimal solutions are nonempty.

Proof. Since (SDP) and (SDD) are strictly feasible, it follows from Corollary 2.1 in p.28 of [10] that the optimal values of (SDP) and (SDD) coincide and their optimal solutions are nonempty. \square

Remark 4.1. Let $A \in S^n$ and $B \in S_{++}^n$ be given. Then, $h_{S_+^n}(A; B) := \min\{t \in \mathbb{R} \mid A \in tB - S_+^n\} = \max\{\text{tr}(AX) \mid \text{tr}(BX) = 1, X \in S_+^n\}$.

From Corollary 4.1, Lemma 4.2, and Remark 4.1, we obtain the following corollary.

Corollary 4.2. Let $f : \mathbb{R}^m \rightarrow S^n$ be a map. Then, exactly one of the following two systems is consistent;

- (i) There exists $x \in \mathbb{R}^m$ such that $f(x) \in -S_{++}^n$;
- (ii) There exists $B \in S_{++}^n$ such that $\max\{\text{tr}(f(x)X) \mid \text{tr}(BX) = 1, X \in S_+^n\} \geq 0$ for all $x \in \mathbb{R}^m$.

This implies a test giving a criterion for satisfaction or dissatisfaction of Slater's condition.

5. CONCLUSION

We have extended some concepts of alternative theorems to set-valued case. Our theorems give a natural generalization to ones having been researched since before. In addition, these result can be used when we are to make sure a set precedes another. Note that there are several ways of reduction of scalarizing functions to ease calculation of their values.

REFERENCES

- [1] J. M. Borwein and A. S. Lewis, *Convex analysis and nonlinear optimization*, Springer, New York, 2000.
- [2] P. Gordan, *Über die Auflösung linearer Gleichungen mit reellen coefficienten*, Mathematische Annalen, **6** (1873) 23–28.
- [3] A. Göpfert, H. Riahi, C. Tammer, and C. Zălinescu, *Variational methods in partially ordered spaces*, Springer, New York, 2003.
- [4] V. Jeyakumar, *A generalization of a minimax theorem of Fan via a theorem of the alternative*, J. Optim. Theory Appl. **48** (1986) 525–533.

- [5] D. Kuroiwa, T. Tanaka, and T. X. D. Ha, *On convexity of set-valued maps*, Nonlinear Anal. **30** (1997), 1487–1496.
- [6] I. Kuwano, T. Tanaka, and S. Yamada, *Characterization of nonlinear scalarizing function for set-valued maps*, in Proceedings of the Asian Conference on Nonlinear Analysis and Optimization, S. Akashi, W. Takahashi, and T. Tanaka (eds.), Yokohama Publishers, Yokohama, 2009, pp.193–204.
- [7] Z. Li, *A theorem of the alternative and its application to the optimization of set-valued maps*, J. Optim. Theory Appl. **100** (1999) 365–375.
- [8] S. Nishizawa, M. Onoduka, and T. Tanaka, *Alternative theorems for set-valued maps based on a nonlinear scalarization*, Pacific J. Optim. **1** (2005) 147–159.
- [9] X. M. Yang, X. Q. Yang, and G. Y. Chen, *Theorems of the alternative and optimization with set-valued maps*, J. Optim. Theory Appl. **107** (2000) 627–640.
- [10] E. de Klerk: *Aspects of semidefinite programming: interior point algorithms and selected applications*, Kluwer Academic Publishers, Dordrecht (2002).

GRADUATE SCHOOL OF SCIENCE AND TECHNOLOGY, NIIGATA UNIVERSITY, NIIGATA, JAPAN

E-mail address (Yuto Ogata): y-ogata@m.sc.niigata-u.ac.jp

E-mail address (Yutaka Saito): ysaito@m.sc.niigata-u.ac.jp

E-mail address (Tamaki Tanaka): tamaki@math.sc.niigata-u.ac.jp

新潟大学大学院自然科学研究科 小形優人, 齋藤裕, 田中 環

DEPARTMENT OF APPLIED MATHEMATICS, PUKYONG NATIONAL UNIVERSITY, BUSAN, KOREA

E-mail address (Gue Myung Lee): gmllee@pknu.ac.kr

E-mail address (Jae Hyoun Lee): mc7558@naver.com

釜慶大学校応用数学科 GUE MYUNG LEE, JAE HYOUNG LEE